




Meaningful Stories. Meaningful Hope.



Chris
Actual LIBTAYO Patient

Finding the Right Doctor or Treatment Center for You

What I've Learned About Taking Charge of My Advanced CSCC: Three Tips for Becoming Your Own Advocate

Throughout my life, I've always been a planner. Once I decided on a path, I stuck with it. But when I was diagnosed with metastatic cutaneous squamous cell carcinoma (CSCC), or CSCC that has spread, I had to learn to take a more proactive approach to my healthcare and working with my doctor. I learned that sticking to my initial plan might not always be the right option for me.

It was an adjustment for me, but I learned some valuable lessons from living with CSCC. Here are my tips on how to advocate for yourself.

What is LIBTAYO?

LIBTAYO is a prescription medicine used to treat adults with a type of skin cancer called cutaneous squamous cell carcinoma (CSCC) that has spread or cannot be cured by surgery or radiation.

It is not known if LIBTAYO is safe and effective in children.

Tip 1. Always ask questions.

When I first noticed a lump under my arm, I was initially diagnosed with breast cancer and scheduled chemotherapy.

At the end of my first round of treatment, I went into the exam room to talk with the doctor. My oncologist broke the news that I didn't have breast cancer as they originally thought.

Based on the results of a prior biopsy, and what my tumor looked like now, I had metastatic CSCC or CSCC that has spread. At every step so far, my treatment was supposed to give hope that my condition gets better. And it didn't. I wish I had kept asking questions.

Important Safety Information

What is the most important information I should know about LIBTAYO?

LIBTAYO is a medicine that may treat certain cancers by working with your immune system. LIBTAYO can cause your immune system to attack normal organs and tissues

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Chris
Actual LIBTAYO Patient

Important Safety Information (continued)

in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Tip 2. Do your research. Continue asking until you understand.

You have to take advanced CSCC seriously. Sometimes you will have to do your own research and find the right path forward for you.

Eventually, my oncologist told me that my cancer was not responding to treatment. I felt like my plans for life just shattered all over again. In nine months, I had gone from having breast cancer to metastatic CSCC, to potentially dying. That was harder than anything to wrap my head around.

However, at that point, my oncologist told me there may be another option for me, an immunotherapy called LIBTAYO[®] (cemiplimab-rwlc). LIBTAYO is a prescription medicine used to treat people with a type of skin cancer called cutaneous squamous cell carcinoma (CSCC) that has spread or cannot be cured by surgery or radiation. We discussed the risks and benefits. After that appointment, I came home and did my research. I didn't want to get my hopes up after all I'd been through. By then I had learned to ask more questions and do my own research so I had a better understanding of my condition and what I may expect from my treatment.

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including:

- **Lung problems:** cough, shortness of breath, or chest pain
- **Intestinal problems:** diarrhea (loose stools) or more frequent bowel movements than usual, stools that are black, tarry, sticky or have blood or mucus, or severe stomach-area (abdomen) pain or tenderness
- **Liver problems:** yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach-area (abdomen), dark urine (tea colored), or bleeding or bruising more easily than normal

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Chris
Actual LIBTAYO Patient

Important Safety Information (continued)

- **Hormone gland problems:** headache that will not go away or unusual headaches, eye sensitivity to light, eye problems, rapid heartbeat, increased sweating, extreme tiredness, weight gain or weight loss, feeling more hungry or thirsty than usual, urinating more often than usual, hair loss, feeling cold, constipation, your voice gets deeper, dizziness or fainting, or changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness
- **Kidney problems:** decrease in your amount of urine, blood in your urine, swelling of your ankles, or loss of appetite

Tip 3: Most importantly, search for a team you work well with.

My doctor explained immunotherapy to me—what it was and how it worked and what to possibly expect. The team focused on developing a holistic treatment plan. I felt comfortable with my treatment plan and like I was part of making it.

Metastatic CSCC definitely took away my idea of what my life was going to look like. But it also gave me the insight that life doesn't always follow your plans, and that I should do my part to find out what works best for me. I have learned to look at my options and be involved in my health. If I take care of myself, I can live more for the moment.

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including (continued):

- **Skin problems:** rash, itching, skin blistering or peeling, painful sores or ulcers in mouth or nose, throat, or genital area, fever or flu-like symptoms, or swollen lymph nodes
- **Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with LIBTAYO. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:** chest pain, irregular heartbeat, shortness of breath or swelling of ankles, confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs, double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight, persistent or severe muscle pain or weakness, muscle cramps, low red blood cells, or bruising

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Chris
Actual LIBTAYO Patient

Important Safety Information (continued)

- **Infusion reactions that can sometimes be severe or life-threatening.** Signs and symptoms of infusion reactions may include: nausea, vomiting, chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feel like passing out, fever, back or neck pain, or facial swelling
- **Rejection of a transplanted organ or tissue.** Your healthcare provider should tell you what signs and symptoms you should report and monitor you, depending on the type of organ or tissue transplant that you have had

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including (continued):

- **Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with LIBTAYO. Your healthcare provider will monitor you for these complications

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with LIBTAYO. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with LIBTAYO if you have severe side effects.

Before you receive LIBTAYO, tell your healthcare provider about all your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ or tissue transplant, including corneal transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Chris
Actual LIBTAYO Patient

Important Safety Information (continued)

- are pregnant or plan to become pregnant. LIBTAYO can harm your unborn baby
Females who are able to become pregnant:
 - Your healthcare provider will give you a pregnancy test before you start treatment
 - You should use an effective method of birth control during your treatment and for at least 4 months after your last dose of LIBTAYO. Talk to your healthcare provider about birth control methods that you can use during this time
 - Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with LIBTAYO
- are breastfeeding or plan to breastfeed. It is not known if LIBTAYO passes into your breast milk. Do not breastfeed during treatment and for at least 4 months after the last dose of LIBTAYO

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of LIBTAYO when used alone to treat CSCC that has spread or cannot be cured by surgery or radiation, BCC, or NSCLC include tiredness, muscle or bone pain, rash, diarrhea, and low levels of red blood cells (anemia). The most common side effects of LIBTAYO when used alone to help prevent CSCC from coming back include rash and itching. The most common side effects of LIBTAYO when used in combination with platinum-containing chemotherapy to treat NSCLC include hair loss, muscle or bone pain, nausea, tiredness, numbness, pain, tingling, or burning in your hands or feet, and decreased appetite.


These are not all the possible side effects of LIBTAYO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Regeneron Pharmaceuticals at 1-877-542-8296.

Please [click here for full Prescribing Information](#), including [Medication Guide](#).

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Meaningful Stories. Meaningful Hope.



Chris
Actual LIBTAYO Patient

Important Safety Information (continued)

What is LIBTAYO?

LIBTAYO is a prescription medicine used to treat adults with a type of skin cancer called cutaneous squamous cell carcinoma (CSCC) that has spread or cannot be cured by surgery or radiation.

It is not known if LIBTAYO is safe and effective in children.

REGENERON[®]

© 2025 Regeneron Pharmaceuticals, Inc. All rights reserved.
US.LIB.25.09.0172 10/25