

Dave
Actual LIBTAYO Patient

Meaningful Stories. Meaningful Hope.

What is Advanced CSCC?

I have some choice words for advanced cutaneous squamous cell carcinoma (CSCC), but I can tell you that it's an inconvenience and an ongoing threat to my health. It's also a condition I have learned to live with. I may not like it. It's certainly not a friend. It's not even a happy acquaintance. But it's a condition that I'm making peace with.

CSCC is the second most common type of skin cancer in the United States. It starts in squamous cells within the surface of the skin and, in advanced stages, invades deeper layers of the skin or spreads to lymph nodes (which may become swollen, painful, or tender) or other parts of the body. If CSCC grows around or into the nerves, it can also cause numbness and pain, as well as muscle weakness. Surgery and/or radiation may be used to remove CSCC tumors on the head, neck, and other parts of the body. However, in advanced stages, certain patients may not be able to be cured with surgery or radiation.

Of course, advanced CSCC can look and feel different from person to person. Everyone has different areas that are affected and different problems and issues to deal with.

What is LIBTAYO?

LIBTAYO is a prescription medicine used to treat adults with a type of skin cancer called cutaneous squamous cell carcinoma (CSCC) that has spread or cannot be cured by surgery or radiation.


It is not known if LIBTAYO is safe and effective in children.

Important Safety Information

What is the most important information I should know about LIBTAYO?

LIBTAYO is a medicine that may treat certain cancers by working with your immune system. LIBTAYO can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Dave
Actual LIBTAYO Patient

Meaningful Stories. Meaningful Hope.

Below the surface

With my most recent CSCC, it started in the lower layers of the skin. On the surface, there wasn't a typical lesion or roughness. The skin looked normal. There wasn't redness. It was thick and hard, but otherwise, it didn't look bad. Then it got thicker.

I'd never had anything like that before and didn't know what to make of it. So of course, I went to my dermatologist who referred me to a surgeon and an oncologist. The team told me that my CSCC had become advanced, which meant that it had spread into deep layers of the skin. I'm fortunate that I was referred to an oncologist who knew about LIBTAYO as a treatment option for CSCC that has spread or cannot be cured by surgery or radiation.


Living with advanced CSCC has not been easy, and I stay vigilant. I get frequent checkups and work with my treatment team.

Important Safety Information (continued)

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including:

- **Lung problems:** cough, shortness of breath, or chest pain
- **Intestinal problems:** diarrhea (loose stools) or more frequent bowel movements than usual, stools that are black, tarry, sticky or have blood or mucus, or severe stomach-area (abdomen) pain or tenderness
- **Liver problems:** yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach-area (abdomen), dark urine (tea colored), or bleeding or bruising more easily than normal
- **Hormone gland problems:** headache that will not go away or unusual headaches, eye sensitivity to light, eye problems, rapid heartbeat, increased sweating, extreme tiredness, weight gain or weight loss, feeling more hungry or thirsty than usual, urinating more often than usual, hair loss, feeling cold, constipation, your voice gets deeper, dizziness or fainting, or changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness
- **Kidney problems:** decrease in your amount of urine, blood in your urine, swelling of your ankles, or loss of appetite
- **Skin problems:** rash, itching, skin blistering or peeling, painful sores or ulcers in mouth or nose, throat, or genital area, fever or flu-like symptoms, or swollen lymph nodes

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Dave
Actual LIBTAYO Patient

Meaningful Stories. Meaningful Hope.

Important Safety Information (continued)

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including (continued):


- **Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with LIBTAYO. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:** chest pain, irregular heartbeat, shortness of breath or swelling of ankles, confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs, double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight, persistent or severe muscle pain or weakness, muscle cramps, low red blood cells, or bruising
- **Infusion reactions that can sometimes be severe or life-threatening.** Signs and symptoms of infusion reactions may include: nausea, vomiting, chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feel like passing out, fever, back or neck pain, or facial swelling
- **Rejection of a transplanted organ or tissue.** Your healthcare provider should tell you what signs and symptoms you should report and monitor you, depending on the type of organ or tissue transplant that you have had
- **Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with LIBTAYO. Your healthcare provider will monitor you for these complications

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with LIBTAYO. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with LIBTAYO if you have severe side effects.

Before you receive LIBTAYO, tell your healthcare provider about all your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ or tissue transplant, including corneal transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Dave
Actual LIBTAYO Patient

Meaningful Stories. Meaningful Hope.

Important Safety Information (continued)

Before you receive LIBTAYO, tell your healthcare provider about all your medical conditions, including if you (continued):

- have received radiation treatment to your chest area
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. LIBTAYO can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider will give you a pregnancy test before you start treatment
- You should use an effective method of birth control during your treatment and for at least 4 months after your last dose of LIBTAYO. Talk to your healthcare provider about birth control methods that you can use during this time
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with LIBTAYO
- are breastfeeding or plan to breastfeed. It is not known if LIBTAYO passes into your breast milk. Do not breastfeed during treatment and for at least 4 months after the last dose of LIBTAYO


Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of LIBTAYO when used alone to treat CSCC that has spread or cannot be cured by surgery or radiation, BCC, or NSCLC include tiredness, muscle or bone pain, rash, diarrhea, and low levels of red blood cells (anemia). The most common side effects of LIBTAYO when used alone to help prevent CSCC from coming back include rash and itching. The most common side effects of LIBTAYO when used in combination with platinum-containing chemotherapy to treat NSCLC include hair loss, muscle or bone pain, nausea, tiredness, numbness, pain, tingling, or burning in your hands or feet, and decreased appetite.

These are not all the possible side effects of LIBTAYO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Regeneron Pharmaceuticals at 1-877-542-8296.

Please [click here for full Prescribing Information](#), including [Medication Guide](#).

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).



Dave
Actual LIBTAYO Patient

Meaningful Stories. Meaningful Hope.

What is LIBTAYO?

LIBTAYO is a prescription medicine used to treat adults with a type of skin cancer called cutaneous squamous cell carcinoma (CSCC) that has spread or cannot be cured by surgery or radiation.

It is not known if LIBTAYO is safe and effective in children.

REGENERON[®]

© 2025 Regeneron Pharmaceuticals, Inc. All rights reserved.
US.LIB.25.09.0176 10/25

Please see additional Important Safety Information throughout and [click here for full Prescribing Information](#), including [Medication Guide](#).